Euro. Phys. J. C **72**, 1907 (2012)

arXiv:1112.3700v2 [hep-ph] 28 Mar 2012

Extracting Rephase-invariant CP and CPT Violating Parameters from Asymmetries of Time-Ordered Integrated Rates of Correlated Decays of Entangled Mesons

Zhijie Huang and Yu Shi*
Department of Physics, Fudan University, Shanghai, 200433, China

We present a general model-independent formalism of measuring CP and CPT violating parameters through time-ordered integrated rates of correlated decays of $C = \pm 1$ entangled states of neutral pseudoscalar mesons. We give the general formulae of CP and CPT violating parameters in terms of four measurable asymmetries defined for the time-ordered integrated rates, applicable to all kinds of decay product. Two special cases which are often realized in experiments are discussed specifically.

PACS numbers: 11.30.Er, 13.20.-v, 14.40.-n, 03.65.Ud

---

* yushi@fudan.edu.cn

I. INTRODUCTION

Precise measurement of CP violating parameters in weak decays is an active topic. On the other hand, testing CPT symmetry is also of great importance, as some physics beyond the standard model may lead to its breaking. Nowadays, neutral K and B mesons are routinely produced as quantum-entangled or EPR-correlated pairs in ϕ and B factories [1–4]. Hence it is highly interesting to explore their use in examining CP and CPT symmetries [1–3, 5–10], and in measuring relevant parameters [11, 12]. Especially, systematic proposals were made on exploring CP and CPT violating parameters in correlated decays of meson pairs in $C=-1$ state, by using decay rates and their asymmetries [6, 7]. In this paper, we present a general and rigorous calculation of time-ordered integrated rates of the correlated decays of $C=\pm 1$ entangled meson pairs, taking into account both CP violation and possible CPT violation. Subsequently CP and CPT violating parameters are expressed as functions of the four asymmetries defined for the time-ordered integrated rates of the correlated decays of the two entangled states. Our result provides a general method of measuring rephase-invariant CP and CPT violating parameters in correlated decays of entangled mesons into any kind of product. This extends a rephase-invariant formalism of CP and CPT violating observables [13] to $C=\pm 1$ entangled mesons.

In Sec. II, we review the single-meson decays using two complex numbers parameterizing indirect CP and CPT violations. In Sec. III, we consider the entangled states $|\psi_{\pm}\rangle$ and calculate the general expressions of the decay rates as functions of four single-meson decay amplitudes. In Sec. IV, we integrate the decay rates over all times with two time orders, thereby obtain four time-ordered integrated rates. We give both exact results and approximate expressions up to the first order of the CP and CPT violating parameters. Subsequently, we obtaining the four real parameters as functions of the four asymmetries defined for the four time-ordered integrated rates. In Sec. V, two special cases are treated for which the expressions are much simpler than the general expressions in Sec. IV. Discussions and a summary are made in Sec. VI.

II. REVIEW OF CP AND CPT VIOLATING PARAMETERS IN SINGLE-MESON DECAYS

An arbitrary state of a pseudoscalar meson can be written as

$$|M(t)\rangle = \psi(t)|M^0\rangle + \bar{\psi}(t)|\bar{M}^0\rangle, \tag{1}$$

where $|M^0\rangle$ is the pseudoscalar meson state and $|\bar{M}^0\rangle$ is its antiparticle state, $\psi(t)$ and $\bar{\psi}(t)$ are superposition coefficients. In Wigner-Weisskopf approximation, the time evolution of $|M(t)\rangle$ is described as

$$i\frac{d}{dt}\begin{pmatrix}\psi(t)\\ \bar{\psi}(t)\end{pmatrix} = \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22}\end{pmatrix}\begin{pmatrix}\psi(t)\\ \bar{\psi}(t)\end{pmatrix}, \tag{2}$$

where $H_{ij}=M_{ij}-i\Gamma_{ij}/2$, with the dispersive part M_{ij} and the absorptive part Γ_{ij} are hermitian. Define [13]

$$\delta_M \equiv \frac{H_{22}-H_{11}}{\sqrt{H_{12}H_{21}}}, \tag{3}$$

$$\frac{q}{p} \equiv \sqrt{\frac{H_{21}}{H_{12}}} \equiv \frac{1-\varepsilon_M}{1+\varepsilon_M}. \tag{4}$$

The eigenstates of the Hamiltonian are

$$\begin{aligned} |M_1\rangle &= p_1|M^0\rangle + q_1|\bar{M}^0\rangle, \\ |M_2\rangle &= p_2|M^0\rangle - q_2|\bar{M}^0\rangle, \end{aligned} \tag{5}$$

with the respective eigenvalues

$$\begin{aligned} \lambda_1 &= H_{11} + \sqrt{H_{12}H_{21}}\left(\frac{\delta_M}{2} + \sqrt{1+\frac{\delta_M^2}{4}}\right), \\ \lambda_2 &= H_{22} - \sqrt{H_{12}H_{21}}\left(\frac{\delta_M}{2} + \sqrt{1+\frac{\delta_M^2}{4}}\right). \end{aligned} \tag{6}$$

Therefore we obtain

$$\begin{aligned} |M^0\rangle &= \frac{q_2|M_1\rangle + q_1|M_2\rangle}{p_1q_2+p_2q_1}, \\ |\bar{M}^0\rangle &= \frac{p_2|M_1\rangle - p_1|M_2\rangle}{p_1q_2+p_2q_1}. \end{aligned} \tag{7}$$

For a single meson whose initial state is $|M^0\rangle$, we have

$$|M^0(t)\rangle = \frac{p_1 q_2 e^{-i\lambda_1 t} + p_2 q_1 e^{-i\lambda_2 t}}{p_1 q_2 + p_2 q_1}|M_0\rangle + \frac{q_1 q_2 (e^{-i\lambda_1 t} - e^{-i\lambda_2 t})}{p_1 q_2 + p_2 q_1}|\bar{M}_0\rangle. \tag{8}$$

Similarly, for a single meson whose initial state is $|\bar{M}^0\rangle$,

$$|\bar{M}^0(t)\rangle = \frac{p_1 p_2 (e^{-i\lambda_1 t} - e^{-i\lambda_2 t})}{p_1 q_2 + p_2 q_1}|M_0\rangle + \frac{p_2 q_1 e^{-i\lambda_1 t} + p_1 q_2 e^{-i\lambda_2 t}}{p_1 q_2 + p_2 q_1}|\bar{M}_0\rangle. \tag{9}$$

Define

$$\frac{1+\Delta_M}{1-\Delta_M} \equiv \frac{\delta_M}{2} + \sqrt{1+\frac{\delta_M^2}{4}}, \tag{10}$$

then [13]

$$\frac{p_1}{q_1} = \frac{p}{q}\frac{1-\Delta_M}{1+\Delta_M}, \tag{11}$$

$$\frac{p_2}{q_2} = \frac{p}{q}\frac{1+\Delta_M}{1-\Delta_M}. \tag{12}$$

Therefore

$$p_1 \propto \frac{1}{\sqrt{2}}(1+\varepsilon_M)(1-\Delta_M), \tag{13}$$

$$q_1 \propto \frac{1}{\sqrt{2}}(1-\varepsilon_M)(1+\Delta_M), \tag{14}$$

$$p_2 \propto \frac{1}{\sqrt{2}}(1+\varepsilon_M)(1+\Delta_M), \tag{15}$$

$$q_2 \propto \frac{1}{\sqrt{2}}(1-\varepsilon_M)(1-\Delta_M). \tag{16}$$

Therefore, up to the order of $O(\Delta_M)$ and $O(\varepsilon_M)$, we can simplify Equations (8) and (9) as

$$|M^0(t)\rangle = [g_+(t) - 2\Delta_M g_-(t)]|M_0\rangle + (1+2\varepsilon_M)g_-(t)|\bar{M}_0\rangle, \tag{17}$$

$$|\bar{M}^0(t)\rangle = (1-2\varepsilon_M)g_-(t)|M_0\rangle + [g_+(t) + 2\Delta_M g_-(t)]|\bar{M}_0\rangle, \tag{18}$$

where

$$g_+ \equiv \frac{1}{2}(e^{-i\lambda_1 t} + e^{-i\lambda_2 t}), \tag{19}$$

$$g_- \equiv \frac{1}{2}(e^{-i\lambda_1 t} - e^{-i\lambda_2 t}). \tag{20}$$

If CPT is conserved, then $H_{11} = H_{22}$, therefore $\Delta_M = 0$. Independent of the situation of CPT, if CP is conserved, then $\varepsilon_M = 0$. If T is conserved, then $M_{12}^*/M_{12} = \Gamma_{12}^*/\Gamma_{12}$, therefore $\Re\varepsilon_M = 0$. If CPT and CP are both conserved, we have T conservation. But T conservation can also be valid without requiring the conservation of CPT and CP. T conservation together with CP violation means $\Im\varepsilon_M \neq 0$.

III. CORRELATED DECAYS OF ENTANGLED STATES

Suppose the initial state of two mesons a and b is the entangled state of $C = \pm 1$,

$$|\psi_\pm\rangle = \frac{1}{\sqrt{2}}[|M_0\rangle_a|\bar{M}_0\rangle_b \pm |\bar{M}_0\rangle_a|M_0\rangle_b]. \tag{21}$$

After time t_a, the final state of a is $|f_a\rangle$, while after time t_b, the final state of b is $|f_b\rangle$.

Define the decay amplitudes of M_0 and $\bar{M}_0$,

$$g_1 \equiv \langle f_a | M_0 \rangle, \tag{22}$$

$$g_2 \equiv \langle f_b | M_0 \rangle, \tag{23}$$

$$h_1 \equiv \langle f_a | \bar{M}_0 \rangle, \tag{24}$$

$$h_2 \equiv \langle f_b | \bar{M}_0 \rangle. \tag{25}$$

The decay amplitude of the entangled meson pair can be obtained from (17) and (18) as

$$A^{\pm}_{f_a f_b}(t_a, t_b) = \frac{1}{\sqrt{2}} \big[\langle f_a | M_0(t_a) \rangle \langle f_b | \bar{M}_0(t_b) \rangle \pm \langle f_a | \bar{M}_0(t_a) \rangle \langle f_b | M_0(t_b) \rangle \big]. \tag{26}$$

The decay rates for the entangled states are thus

$$\begin{aligned}
\Gamma^{+}_{f_a f_b}(t_a, t_b) &= |A^{+}_{f_a f_b}(t_a, t_b)|^2 \\
&= \frac{1}{2} e^{-\Gamma_1 (t_a + t_b)} |\alpha_1|^2 + \frac{1}{2} e^{-\Gamma_2 (t_a + t_b)} |\alpha_2|^2 + \frac{1}{2} e^{-\Gamma_1 t_a - \Gamma_2 t_b} |\alpha_3|^2 + \frac{1}{2} e^{-\Gamma_2 t_a - \Gamma_1 t_b} |\alpha_4|^2 \\
&\quad + e^{-\Gamma (t_a + t_b)} \Re[\alpha_1^* \alpha_2 e^{-i \Delta m (t_a + t_b)}] + e^{-\Gamma_1 t_a - \Gamma t_b} \Re[\alpha_1^* \alpha_3 e^{-i \Delta m t_b}] \\
&\quad + e^{-\Gamma t_a - \Gamma_1 t_b} \Re[\alpha_1^* \alpha_4 e^{-i \Delta m t_a}] + e^{-\Gamma t_a - \Gamma_2 t_b} \Re[\alpha_2^* \alpha_3 e^{i \Delta m t_a}] \\
&\quad + e^{-\Gamma_2 t_a - \Gamma t_b} \Re[\alpha_2^* \alpha_4 e^{i \Delta m t_b}] + e^{-\Gamma (t_a + t_b)} \Re[\alpha_3^* \alpha_4 e^{i \Delta m (t_b - t_a)}],
\end{aligned} \tag{27}$$

$$\begin{aligned}
\Gamma^{-}_{f_a f_b}(t_a, t_b) &= |A^{-}_{f_a f_b}(t_a, t_b)|^2 \\
&= \frac{1}{2} e^{-\Gamma_1 (t_a + t_b)} |\beta_1|^2 + \frac{1}{2} e^{-\Gamma_2 (t_a + t_b)} |\beta_2|^2 + \frac{1}{2} e^{-\Gamma_1 t_a - \Gamma_2 t_b} |\beta_3|^2 + \frac{1}{2} e^{-\Gamma_2 t_a - \Gamma_1 t_b} |\beta_4|^2 \\
&\quad + e^{-\Gamma (t_a + t_b)} \Re[\beta_1^* \beta_2 e^{-i \Delta m (t_a + t_b)}] + e^{-\Gamma_1 t_a - \Gamma t_b} \Re[\beta_1^* \beta_3 e^{-i \Delta m t_b}] \\
&\quad + e^{-\Gamma t_a - \Gamma_1 t_b} \Re[\beta_1^* \beta_4 e^{-i \Delta m t_a}] + e^{-\Gamma t_a - \Gamma_2 t_b} \Re[\beta_2^* \beta_3 e^{i \Delta m t_a}] \\
&\quad + e^{-\Gamma_2 t_a - \Gamma t_b} \Re[\beta_2^* \beta_4 e^{i \Delta m t_b}] + e^{-\Gamma (t_a + t_b)} \Re[\beta_3^* \beta_4 e^{i \Delta m (t_b - t_a)}],
\end{aligned} \tag{28}$$

where

$$\begin{aligned}
\Delta m &\equiv m_2 - m_1, \\
\Gamma &\equiv \frac{\Gamma_1 + \Gamma_2}{2}, \\
\Delta \Gamma &\equiv \Gamma_2 - \Gamma_1, \\
\alpha_1 &= \frac{1}{2}(g_1 + h_1)(g_2 + h_2) + (\Delta_M + \varepsilon_M)(h_1 h_2 - g_1 g_2) - (\Delta_M^2 + \varepsilon_M^2)(g_1 h_2 + g_2 h_1) \\
&\quad + 2 \Delta_M \varepsilon_M (g_1 g_2 + h_1 h_2), \\
\alpha_2 &= -\frac{1}{2}(g_1 - h_1)(g_2 - h_2) + (\Delta_M - \varepsilon_M)(h_1 h_2 - g_1 g_2) - (\Delta_M^2 + \varepsilon_M^2)(g_1 h_2 + g_2 h_1) \\
&\quad + 2 \Delta_M \varepsilon_M (g_1 g_2 + h_1 h_2), \\
\alpha_3 &= \Delta_M (g_1 + h_1)(g_2 - h_2) + (\Delta_M^2 + \varepsilon_M^2)(g_1 h_2 + g_2 h_1) - 2 \Delta_M \varepsilon_M (g_1 g_2 + h_1 h_2), \\
\alpha_4 &= \Delta_M (g_1 - h_1)(g_2 + h_2) + (\Delta_M^2 + \varepsilon_M^2)(g_1 h_2 + g_2 h_1) - 2 \Delta_M \varepsilon_M (g_1 g_2 + h_1 h_2), \\
\beta_1 &= \beta_2 = (\Delta_M^2 - \varepsilon_M^2)(g_2 h_1 - g_1 h_2), \\
\beta_3 &= \frac{1}{2}(g_1 + h_1)(h_2 - g_2) - \Delta_M (g_2 h_1 + g_1 h_2) + \varepsilon_M (g_1 g_2 + h_1 h_2) + (\Delta_M^2 - \varepsilon_M^2)(g_1 h_2 - g_2 h_1), \\
\beta_4 &= \frac{1}{2}(g_1 - h_1)(g_2 + h_2) + \Delta_M (g_2 h_1 + g_1 h_2) - \varepsilon_M (g_1 g_2 + h_1 h_2) + (\Delta_M^2 - \varepsilon_M^2)(g_1 h_2 - g_2 h_1).
\end{aligned}$$

IV. TIME-ORDERED INTEGRATED DECAY RATES

Now consider two kinds of decay of an entangled state $|\psi_{\pm}\rangle$. One, whose integrated rate is denoted as $R^{\pm}_{ba}$, is that the decay of b into f_b precedes that of a into f_a, i.e. $t_b \le t_a$. The other, whose integrated rate is denoted as $R^{\pm}_{ab}$, is of the inverse time ordering, i.e. $t_b \geq t_a$. That is,

$$R_{ba}^{\pm} = \int_0^{\infty} dt_a \int_0^{t_a} dt_b \Gamma^{\pm}_{f_a f_b}(t_a, t_b), \tag{29}$$

$$R_{ab}^{\pm} = \int_0^{\infty} dt_a \int_{t_a}^{\infty} dt_b \Gamma^{\pm}_{f_a f_b}(t_a, t_b). \tag{30}$$

Making use of Equations (27) and (28), we obtain

$$\begin{aligned} R_{ba}^{+} &= \frac{|\alpha_1|^2}{4\Gamma_1^2} + \frac{|\alpha_2|^2}{4\Gamma_2^2} + \frac{|\alpha_3|^2}{4\Gamma_1\Gamma} + \frac{|\alpha_4|^2}{4\Gamma_2\Gamma} + \Re[\frac{\alpha_1^*\alpha_2}{2(\Gamma+i\Delta m)^2}] + \Re[\frac{\alpha_1^*\alpha_3}{\Gamma_1(\Gamma_1+\Gamma+i\Delta m)}] \\ &+\Re[\frac{\alpha_1^*\alpha_4}{(\Gamma+i\Delta m)(\Gamma_1+\Gamma+i\Delta m)}] + \Re[\frac{\alpha_2^*\alpha_3}{(\Gamma-i\Delta m)(\Gamma_2+\Gamma-i\Delta m)}] \\ &+\Re[\frac{\alpha_2^*\alpha_4}{\Gamma_2(\Gamma_2+\Gamma-i\Delta m)}] + \Re[\frac{\alpha_3^*\alpha_4}{2\Gamma(\Gamma+i\Delta m)}] \\ &\simeq A_0 + A_1\Re\Delta_M + A_2\Im\Delta_M - A_3\Re\varepsilon_M - A_4\Im\varepsilon_M, \end{aligned} \tag{31}$$

$$\begin{aligned} R_{ba}^{-} &= \frac{|\beta_1|^2}{4\Gamma_1^2} + \frac{|\beta_2|^2}{4\Gamma_2^2} + \frac{|\beta_3|^2}{4\Gamma_1\Gamma} + \frac{|\beta_4|^2}{4\Gamma_2\Gamma} + \Re[\frac{\beta_1^*\beta_2}{2(\Gamma+i\Delta m)^2}] + \Re[\frac{\beta_1^*\beta_3}{\Gamma_1(\Gamma_1+\Gamma+i\Delta m)}] \\ &+\Re[\frac{\beta_1^*\beta_4}{(\Gamma+i\Delta m)(\Gamma_1+\Gamma+i\Delta m)}] + \Re[\frac{\beta_2^*\beta_3}{(\Gamma-i\Delta m)(\Gamma_2+\Gamma-i\Delta m)}] \\ &+\Re[\frac{\beta_2^*\beta_4}{\Gamma_2(\Gamma_2+\Gamma-i\Delta m)}] + \Re[\frac{\beta_3^*\beta_4}{2\Gamma(\Gamma+i\Delta m)}] \\ &\simeq B_0 + B_1\Re\Delta_M + B_2\Im\Delta_M - B_3\Re\varepsilon_M - B_4\Im\varepsilon_M, \end{aligned} \tag{32}$$

$$\begin{aligned} R_{ab}^{+} &= \frac{|\alpha_1|^2}{4\Gamma_1^2} + \frac{|\alpha_2|^2}{4\Gamma_2^2} + \frac{|\alpha_3|^2}{4\Gamma_2\Gamma} + \frac{|\alpha_4|^2}{4\Gamma_1\Gamma} + \Re[\frac{\alpha_1^*\alpha_2}{2(\Gamma+i\Delta m)^2}] + \Re[\frac{\alpha_1^*\alpha_3}{(\Gamma+i\Delta m)(\Gamma_1+\Gamma+i\Delta m)}] \\ &+\Re[\frac{\alpha_1^*\alpha_4}{\Gamma_1(\Gamma_1+\Gamma+i\Delta m)}] + \Re[\frac{\alpha_2^*\alpha_3}{\Gamma_2(\Gamma_2+\Gamma-i\Delta m)}] \\ &+\Re[\frac{\alpha_2^*\alpha_4}{(\Gamma-i\Delta m)(\Gamma_2+\Gamma-i\Delta m)}] + \Re[\frac{\alpha_3^*\alpha_4}{2\Gamma(\Gamma-i\Delta m)}] \\ &\simeq C_0 + C_1\Re\Delta_M + C_2\Im\Delta_M - C_3\Re\varepsilon_M - C_4\Im\varepsilon_M, \end{aligned} \tag{33}$$

$$\begin{aligned} R_{ab}^{-} &= \frac{|\beta_1|^2}{4\Gamma_1^2} + \frac{|\beta_2|^2}{4\Gamma_2^2} + \frac{|\beta_3|^2}{4\Gamma_2\Gamma} + \frac{|\beta_4|^2}{4\Gamma_1\Gamma} + \Re[\frac{\beta_1^*\beta_2}{2(\Gamma+i\Delta m)^2}] + \Re[\frac{\beta_1^*\beta_3}{(\Gamma+i\Delta m)(\Gamma_1+\Gamma+i\Delta m)}] \\ &+\Re[\frac{\beta_1^*\beta_4}{\Gamma_1(\Gamma_1+\Gamma+i\Delta m)}] + \Re[\frac{\beta_2^*\beta_3}{\Gamma_2(\Gamma_2+\Gamma-i\Delta m)}] \\ &+\Re[\frac{\beta_2^*\beta_4}{(\Gamma-i\Delta m)(\Gamma_2+\Gamma-i\Delta m)}] + \Re[\frac{\beta_3^*\beta_4}{2\Gamma(\Gamma-i\Delta m)}] \\ &\simeq D_0 + D_1\Re\Delta_M + D_2\Im\Delta_M - D_3\Re\varepsilon_M - D_4\Im\varepsilon_M, \end{aligned} \tag{34}$$

where $\{A_i\}, \{B_i\}, \{C_i\}$ and $\{D_i\} (i = 0, ..., 4)$ are given in the appendix.

Now we consider the asymmetries defined for these four time-ordered integrated decay rates,

$$R_1 \equiv \frac{R_{ba}^{+} - R_{ab}^{+}}{R_{ba}^{+} + R_{ab}^{+}}, \tag{35}$$

$$R_2 \equiv \frac{R_{ba}^{-} - R_{ab}^{-}}{R_{ba}^{-} + R_{ab}^{-}}, \tag{36}$$

$$R_3 \equiv \frac{R_{ba}^{+} - R_{ba}^{-}}{R_{ba}^{+} + R_{ba}^{-}}, \tag{37}$$

$$R_4 \equiv \frac{R_{ab}^{+} - R_{ab}^{-}}{R_{ab}^{+} + R_{ab}^{-}}, \tag{38}$$

which are calculated to be

$$
\begin{aligned}
R_1 &\simeq \frac{A_1 - C_1}{2A_0}\Re\Delta_M + \frac{A_2 - C_2}{2A_0}\Im\Delta_M, \qquad (39)\\
R_2 &\simeq \frac{B_0 - D_0}{B_0 + D_0} + \frac{2(B_1D_0 - B_0D_1)}{(B_0 + D_0)^2}\Re\Delta_M + \frac{2(B_2D_0 - B_0D_2)}{(B_0 + D_0)^2}\Im\Delta_M\\
&\quad + \frac{2(B_3D_0 - B_0D_3)}{(B_0 + D_0)^2}\Re\varepsilon_M + \frac{2(B_4D_0 - B_0D_4)}{(B_0 + D_0)^2}\Im\varepsilon_M, \qquad (40)\\
R_3 &\simeq \frac{A_0 - B_0}{A_0 + B_0} + \frac{2(A_1B_0 - A_0B_1)}{(A_0 + B_0)^2}\Re\Delta_M + \frac{2(A_2B_0 - A_0B_2)}{(A_0 + B_0)^2}\Im\Delta_M\\
&\quad + \frac{2(A_3B_0 - A_0B_3)}{(A_0 + B_0)^2}\Re\varepsilon_M + \frac{2(A_4B_0 - A_0B_4)}{(A_0 + B_0)^2}\Im\varepsilon_M, \qquad (41)\\
R_4 &\simeq \frac{C_0 - D_0}{C_0 + D_0} + \frac{2(C_1D_0 - C_0D_1)}{(C_0 + D_0)^2}\Re\Delta_M + \frac{2(C_2D_0 - C_0D_2)}{(C_0 + D_0)^2}\Im\Delta_M\\
&\quad + \frac{2(C_3D_0 - C_0D_3)}{(C_0 + D_0)^2}\Re\varepsilon_M + \frac{2(C_4D_0 - C_0D_4)}{(C_0 + D_0)^2}\Im\varepsilon_M, \qquad (42)
\end{aligned}
$$

which can be rewritten as

$$
\begin{pmatrix} R_1 \\ R_2 - \frac{B_0-D_0}{B_0+D_0} \\ R_3 - \frac{A_0-B_0}{A_0+B_0} \\ R_4 - \frac{C_0-D_0}{C_0+D_0} \end{pmatrix} = K \begin{pmatrix} \Re\Delta_M \\ \Im\Delta_M \\ \Re\varepsilon_M \\ \Im\varepsilon_M \end{pmatrix}, \tag{43}
$$

with

$$
K \equiv \begin{pmatrix}
\frac{A_1-C_1}{2A_0} & \frac{A_2-C_2}{2A_0} & 0 & 0 \\
2\frac{B_1D_0-B_0D_1}{(B_0+D_0)^2} & 2\frac{B_2D_0-B_0D_2}{(B_0+D_0)^2} & 2\frac{B_3D_0-B_0D_3}{(B_0+D_0)^2} & 2\frac{B_4D_0-B_0D_4}{(B_0+D_0)^2} \\
2\frac{A_1B_0-A_0B_1}{(A_0+B_0)^2} & 2\frac{A_2B_0-A_0B_2}{(A_0+B_0)^2} & 2\frac{A_3B_0-A_0B_3}{(A_0+B_0)^2} & 2\frac{A_4B_0-A_0B_4}{(A_0+B_0)^2} \\
2\frac{C_1D_0-C_0D_1}{(C_0+D_0)^2} & 2\frac{C_2D_0-C_0D_2}{(C_0+D_0)^2} & 2\frac{C_3D_0-C_0D_3}{(C_0+D_0)^2} & 2\frac{C_4D_0-C_0D_4}{(C_0+D_0)^2}
\end{pmatrix}. \tag{44}
$$

Therefore, CP and CPT violating parameters are obtained as

$$
\begin{pmatrix} \Re\Delta_M \\ \Im\Delta_M \\ \Re\varepsilon_M \\ \Im\varepsilon_M \end{pmatrix} = K^{-1} \begin{pmatrix} R_1 \\ R_2 - \frac{B_0-D_0}{B_0+D_0} \\ R_3 - \frac{A_0-B_0}{A_0+B_0} \\ R_4 - \frac{C_0-D_0}{C_0+D_0} \end{pmatrix}, \tag{45}
$$

where, in terms of the Levi-Civita symbol ϵ_{ijkl}, $\Lambda_{ijk} \equiv (A_iB_0 - A_0B_i)(B_jD_0 - B_0D_j)(C_kD_0 - C_0D_k)$ and $\lambda_{ijk} \equiv (A_2 - C_2)\epsilon_{i2jk} - (A_1 - C_1)\epsilon_{1ijk}(i, j, k = 1, 2, 3, 4)$, the matrix elements of K^{-1} are

$$
\begin{aligned}
(K^{-1})_{11} &= -\frac{2A_0\epsilon_{1ijk}\Lambda_{ijk}}{\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{12} &= \frac{(A_2 - C_2)(\Lambda_{324} - \Lambda_{423})(B_0 + D_0)^2}{2(B_2D_0 - B_0D_2)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{13} &= \frac{(A_2 - C_2)(\Lambda_{234} - \Lambda_{243})(A_0 + B_0)^2}{2(A_2B_0 - A_0B_2)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{14} &= -\frac{(A_2 - C_2)(\Lambda_{432} - \Lambda_{342})(C_0 + D_0)^2}{2(C_2D_0 - C_0D_2)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{21} &= \frac{2A_0\epsilon_{i2jk}\Lambda_{ijk}}{\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{22} &= \frac{(A_1 - C_1)(\Lambda_{314} - \Lambda_{413})(B_0 + D_0)^2}{2(B_1D_0 - B_0D_1)\lambda_{ijk}\Lambda_{ijk}},
\end{aligned}
$$

$$
\begin{aligned}
(K^{-1})_{23} &= -\frac{(A_1-C_1)(\Lambda_{134}-\Lambda_{143})(A_0+B_0)^2}{2(A_1B_0-A_0B_1)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{24} &= \frac{(A_1-C_1)(\Lambda_{431}-\Lambda_{341})(C_0+D_0)^2}{2(C_1D_0-C_0D_1)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{31} &= -\frac{2A_0\epsilon_{ij3k}\Lambda_{ijk}}{\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{32} &= \frac{[(\Lambda_{432}-\Lambda_{234})(A_1-C_1)-(\Lambda_{431}-\Lambda_{134})(A_2-C_2)](B_0+D_0)^2}{2(B_3D_0-B_0D_3)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{33} &= \frac{[(\Lambda_{324}-\Lambda_{342})(A_1-C_1)-(\Lambda_{314}-\Lambda_{341})(A_2-C_2)](A_0+B_0)^2}{2(A_3B_0-A_0B_3)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{34} &= \frac{[(\Lambda_{243}-\Lambda_{423})(A_1-C_1)-(\Lambda_{143}-\Lambda_{413})(A_2-C_2)](C_0+D_0)^2}{2(C_3D_0-C_0D_3)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{41} &= \frac{2A_0\epsilon_{ijk4}\Lambda_{ijk}}{\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{42} &= \frac{[(\Lambda_{243}-\Lambda_{342})(A_1-C_1)-(\Lambda_{143}-\Lambda_{341})(A_2-C_2)](B_0+D_0)^2}{2(B_4D_0-B_0D_4)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{43} &= \frac{[(\Lambda_{432}-\Lambda_{423})(A_1-C_1)-(\Lambda_{431}-\Lambda_{413})(A_2-C_2)](A_0+B_0)^2}{2(A_4B_0-A_0B_4)\lambda_{ijk}\Lambda_{ijk}},\\
(K^{-1})_{44} &= \frac{[(\Lambda_{324}-\Lambda_{234})(A_1-C_1)-(\Lambda_{314}-\Lambda_{134})(A_2-C_2)](C_0+D_0)^2}{2(C_3D_0-C_0D_3)\lambda_{ijk}\Lambda_{ijk}}.
\end{aligned}
$$

In general, CP and CPT violating parameters can be obtained from the four asymmetries of the time-ordered integrated rates of the correlated decays of $CP = \pm 1$ entangled states, using the above formulae. But in some cases, the four asymmetries may be independent of CP and CPT violating parameters, as in one of the special cases discussed below.

V. SPECIAL CASES

We now consider the following two special cases [13], for which the calculations are simplified a lot.

First, we consider the situation that $f_a = f$ while $f_b = \bar{f}$, where f and $\bar{f}$ are mutual CP conjugates and are decay products of M^0 and $\bar{M}^0$ respectively. For example, for $M^0 = B^0$, $f = D^-D_S^+, D^-K^+, \pi^-D_S^+, \pi^-K^+$ while $\bar{f} = D^+D_S^-, D^+K^-, \pi^+D_S^-, \pi^+K^-$, respectively.

In this case, the amplitudes satisfy $g_2 = h_1 = 0$. Up to the order $O(\Delta_M)$ and $O(\varepsilon_M)$, $\{\alpha_i\}, \{\beta_i\}$ are given as

$$\alpha_1 = \alpha_2 = \frac{1}{2}g_1h_2, \tag{46}$$

$$\alpha_3 = -\alpha_4 = -\Delta_M g_1h_2, \tag{47}$$

$$\beta_1 = \beta_2 = 0, \tag{48}$$

$$\beta_3 = \big(\frac{1}{2} - \Delta_M\big)g_1h_2, \tag{49}$$

$$\beta_4 = \big(\frac{1}{2} + \Delta_M\big)g_1h_2. \tag{50}$$

Then we obtain $A_{3,4} = B_{1,2,3,4} = C_{3,4} = D_{1,2,3,4} = 0$. Using the expressions given in the appendix, Eq.(43) is simplified as

$$\begin{pmatrix} R_1 \\ R_2 - \frac{B_0-D_0}{B_0+D_0} \\ R_3 - \frac{A_0-B_0}{A_0+B_0} \\ R_4 - \frac{C_0-D_0}{C_0+D_0} \end{pmatrix} = \begin{pmatrix} \frac{A_1-C_1}{2A_0} & \frac{A_2-C_2}{2A_0} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 2\frac{A_1B_0}{(A_0+B_0)^2} & 2\frac{A_2B_0}{(A_0+B_0)^2} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \Re\Delta_M \\ \Im\Delta_M \\ \Re\varepsilon_M \\ \Im\varepsilon_M \end{pmatrix}. \tag{51}$$

In this case, we cannot obtain the CP violating parameter from the four asymmetries, which are independent of the former. Moreover, $R_2 = \frac{B_0-D_0}{B_0+D_0}$ and $R_4 = \frac{C_0-D_0}{C_0+D_0}$ are independent of both CP and CPT violating parameters, while R_1 and R_3 depend only on the CPT violating parameter, but not on the CP violating parameter.

Now we consider the situation that $f_a = f_b = f$, with $f = \bar{f}$ being a CP eigenstate. For example, $f_a = f_b = \pi^+\pi^-$,$\pi^0\pi^0$. In this case, the decay amplitude satisfies $g_1 = g_2$ and $h_1 = h_2$. Hence the $\{\alpha_i\}$, and $\{\beta_i\}$ can be simplified as

$$\alpha_1 = \frac{1}{2}(g_1 + h_1)^2 + (\Delta_M + \varepsilon_M)(h_1^2 - g_1^2) - 2(\Delta_M^2 + \varepsilon_M^2)g_1 h_1 + 2\Delta_M \varepsilon_M (g_1^2 + h_1^2), \tag{52}$$

$$\alpha_2 = -\frac{1}{2}(g_1 - h_1)^2 + (\Delta_M - \varepsilon_M)(h_1^2 - g_1^2) - 2(\Delta_M^2 + \varepsilon_M^2)g_1 h_1 + 2\Delta_M \varepsilon_M (g_1^2 + h_1^2), \tag{53}$$

$$\alpha_3 = \alpha_4 = \Delta_M(g_1^2 - h_1^2) + 2(\Delta_M^2 + \varepsilon_M^2)g_1 h_1 - 2\Delta_M \varepsilon_M (g_1^2 + h_1^2), \tag{54}$$

$$\beta_1 = \beta_2 = 0, \tag{55}$$

$$\beta_3 = -\beta_4 = \frac{1}{2}(g_1^2 - h_1^2) - 2\Delta_M g_1 h_1 + \varepsilon_M(g_1^2 + h_1^2) - 2\Delta_M g_1 h_1 + \varepsilon_M(g_1^2 + h_1^2). \tag{56}$$

Consequently $R^-_{ba} = R^-_{ab} = 2|\varepsilon_M - \Delta_M|^2 |g_1|^2 \left(\frac{1}{\Gamma_1\Gamma_2} - \frac{1}{\Gamma^2 + \Delta m^2}\right)$, which leads to $R_2 = 0$ exactly, no matter whether CP or CPT is violated.

If CP is conserved, then $g_1 = g_2 = h_1 = h_2$, hence $\alpha_1 = 2g_1^2 - 2g_1^2(\Delta_M - \varepsilon_M)^2$, $\alpha_2 = -\alpha_3 = -\alpha_4 = -2g_1^2(\Delta_M - \varepsilon_M)^2$, $\beta_3 = -\beta_4 = 2g_1^2(\varepsilon_M - \Delta_M)$. Consequently, $R^+_{ba} = R^+_{ab} = \frac{|g_1|^4}{\Gamma_1^2} + O\big((\varepsilon_M - \Delta_M)^2\big)$. Then up to the order $O(\varepsilon_M)$ and $O(\Delta_M)$, we have $R_1 = 0$ and $R_3 = R_4 \approx 1$.

VI. DISCUSSIONS AND SUMMARY

Earlier, extensive discussions were made on $|\psi_-\rangle$ [6, 7]. However, most of these discussions concern the time-order asymmetry for integrated decay rates with a given time difference, rather than R_2, which is the asymmetry for the integrations over all time differences. Only for the case of $f_a = \pi^+\pi^-$ while $f_b = \pi^0\pi^0$, was R_2 calculated, with a nonzero value dependent on the CP violating parameter [6, 7]. This is consistent with our results in Sec. IV. Our results are in given in terms of the decay amplitudes defined for $|M_0\rangle$ and $|\bar{M}_0\rangle$, which are related to the indirect CP violating parameter ϵ'.

In ϕ factory and in $\Upsilon(4s)$ resonance in B factories, the branch ratio of C-even meson pairs is negligibly small [7], thus R^+_{ab} and R^+_{ba} cannot be measured there. However, in $\Upsilon(5s)$ resonance operated in B factories such as CLEO and BELLE, the strong decay products include $B_s\bar{B}_s$, $B_s^*\bar{B}_s$ and $B_s\bar{B}_s^*$. As $B_s^* \to B_s\gamma$, the final states of $B_s^*\bar{B}_s$ and $B_s\bar{B}_s^*$ are $B_s\bar{B}_s$ pair in $|\psi_+\rangle$. It turns out that the branch fraction of $|\psi_-\rangle$ and $|\psi_+\rangle$ are 90% and 10%, respectively [11, 14]. Therefore, $\Gamma^{\pm}_{f_a f_b}$ can both be measured once the correlated pairs are identified. In CPLEAR experiment on entangled kaons, the branch ratio between $|\psi_+\rangle$ and $|\psi_-\rangle$ was as considerable as 0.037 [15]. Moreover, we hope that in future, neutral meson production can start with an initial state of even orbital angular momentum so that pairs in $|\psi_+\rangle$ dominate.

To summarize, we have presented a model-independent formalism of extracting rephase-invariant CP and CPT violating parameters from four asymmetries of time-ordered integrated rates of correlated decays of $C = \pm 1$ entangled states of neutral pseudoscalar mesons. Asymmetries of time-ordered integrated rates are used because they are easy to be obtained from experimental data, without the necessity of measuring the exact times of decays. We give the general formulae applicable to all kinds of decay product. In the special case that M^0 and $\bar{M}^0$ decay to CP conjugates, the asymmetries so defined cannot give information on the CP violating parameter. Then one needs to use other observables, such as the time-ordered integrated rates themselves, or the asymmetries defined for specified differences of decay times. In future work, we will make more phenomenological studies and a more detailed comparison with the previous results [6, 7].

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation of China (Grant No. 10875028).

Appendix A: Detailed expressions of A_i, B_i, C_i and D_i

$$\begin{aligned}
A_0 = {} & \frac{|g_1+h_1|^2|g_2+h_2|^2}{8\Gamma_1^2} + \frac{|g_1-h_1|^2|g_2-h_2|^2}{8\Gamma_2^2} \\
& - \frac{\Gamma^2-\Delta m^2}{4(\Gamma^2+\Delta m^2)^2}\left[(|g_1|^2-|h_1|^2)(|g_2|^2-|h_2|^2) - 4\Im(g_1h_1^*)\Im(g_2h_2^*)\right], \\
A_1 = {} & \frac{1}{2\Gamma_1^2}\left[(|h_1h_2|^2-|g_1g_2|^2) - (|g_1|^2-|h_1|^2)\Re(g_2h_2^*) - (|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
& - \frac{1}{2\Gamma_2^2}\left[|h_1h_2|^2-|g_1g_2|^2 + (|g_1|^2-|h_1|^2)\Re(g_2h_2^*) + (|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
& - \frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}\left[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*) + (|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
& + \frac{\Gamma_1+\Gamma}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2(|g_2|^2-|h_2|^2) + \frac{2\Delta m}{\Gamma_1[(\Gamma+\Gamma_1)^2+\Delta m^2]}|g_1+h_1|^2\Im(g_2h_2^*) \\
& - \frac{\Gamma_2+\Gamma}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2(|g_2|^2-|h_2|^2) - \frac{2\Delta m}{\Gamma_2[(\Gamma+\Gamma_2)^2+\Delta m^2]}|g_1-h_1|^2\Im(g_2h_2^*) \\
& + \frac{\Gamma(\Gamma_1+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2(|g_1|^2-|h_1|^2) \\
& - \frac{\Gamma(\Gamma_2+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2(|g_1|^2-|h_1|^2) \\
& + \frac{2\Delta m(\Gamma_1+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2\Im(g_1h_1^*) \\
& - \frac{2\Delta m(\Gamma_2+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2\Im(g_1h_1^*) \\
& - \frac{4\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}\Im(g_1h_1^*g_2h_2^*), \\
A_2 = {} & \frac{1}{2\Gamma_1^2}\left[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*) + (|g_2|^2+|h_2|^2)\Im(g_1h_1^*) + 2\Im(g_1h_1^*g_2h_2^*)\right] \\
& + \frac{1}{2\Gamma_2^2}\left[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*) + (|g_2|^2+|h_2|^2)\Im(g_1h_1^*) - 2\Im(g_1h_1^*g_2h_2^*)\right] \\
& + \frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}\left[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*) + (|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\right] \\
& + \frac{\Delta m(\Gamma_1+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2(|g_1|^2-|h_1|^2) \\
& + \frac{\Delta m(\Gamma_2+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2(|g_1|^2-|h_1|^2) \\
& - 2\frac{\Gamma(\Gamma_1+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2\Im(g_1h_1^*) \\
& - 2\frac{\Gamma(\Gamma_2+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2\Im(g_1h_1^*)
\end{aligned}$$

$$
\begin{aligned}
&+\frac{\Delta m}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2(|g_2|^2-|h_2|^2)\\
&+\frac{\Delta m}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2(|g_2|^2-|h_2|^2)\\
&-\frac{2(\Gamma+\Gamma_1)}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2\Im(g_2h_2^*)\\
&-\frac{2(\Gamma_2+\Gamma)}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2\Im(g_2h_2^*)\\
&-\frac{2\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}(|g_1g_2|^2-|h_1h_2|^2),\\
A_3 =& \frac{1}{2\Gamma_1}\big[|h_1h_2|^2-|g_1g_2|^2-(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]\\
&+\frac{1}{2\Gamma_2}\big[|h_1h_2|^2-|g_1g_2|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]\\
&+\frac{2\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}\big[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\big]\\
&+\frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}(|g_1g_2|^2-|h_1h_2|^2),\\
A_4 =& \frac{1}{2\Gamma_1}\big[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)+2\Im(g_1h_1^*g_2h_2^*)\big]\\
&-\frac{1}{2\Gamma_2}\big[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)-2\Im(g_1h_1^*g_2h_2^*)\big]\\
&+\frac{2\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}\big[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]\\
&-2\frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}\Im(g_1h_1^*g_2h_2^*),\\
B_0 =& \frac{1}{8\Gamma\Gamma_1}|g_1+h_1|^2|g_2-h_2|^2+\frac{1}{8\Gamma\Gamma_2}|g_1-h_1|^2|g_2+h_2|^2\\
&-\frac{1}{4(\Gamma^2+\Delta m^2)}\big[(|g_1|^2-|h_1|^2)(|g_2|^2-|h_2|^2)+4\Im(g_1h_1^*)\Im(g_2h_2^*)\big]\\
&+\frac{\Delta m}{2\Gamma(\Gamma^2+\Delta m^2)}\big[(|g_1|^2-|h_1|^2)\Im(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Im(g_1h_1^*)\big],\\
B_1 =& \frac{1}{2\Gamma\Gamma_2}\big[|g_1h_2|^2-|g_2h_1|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]\\
&-\frac{1}{2\Gamma\Gamma_1}\big[|g_1h_2|^2-|g_2h_1|^2-(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]\\
&-\frac{1}{\Gamma^2+\Delta m^2}\big[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\big]-\frac{2\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}\Im(g_1h_1^*g_2h_2^*),\\
B_2 =& \frac{1}{2\Gamma\Gamma_2}\big[2\Im(g_1h_1^*g_2^*h_2)+(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\big]\\
&-\frac{1}{2\Gamma\Gamma_1}\big[-2\Im(g_1h_1^*g_2^*h_2)+(|h_1|^2+|g_1|^2)\Im(g_2h_2^*)+(|h_2|^2+|g_2|^2)\Im(g_1h_1^*)\big]
\end{aligned}
$$

$$
\begin{aligned}
&- \frac{1}{\Gamma^2+\Delta m^2}\left[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\right] \\
&+ \frac{\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}(|g_1h_2|^2-|g_2h_1|^2), \\
B_3 &= \frac{1}{2\Gamma\Gamma_2}\left[|g_1g_2|^2-|h_1h_2|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_2h_2^*)\right] \\
&+ \frac{1}{2\Gamma\Gamma_1}\left[|h_1h_2|^2-|g_1g_2|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_2h_2^*)\right] \\
&- \frac{\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}\left[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)-(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\right] \\
&+ \frac{1}{\Gamma^2+\Delta m^2}(|g_1g_2|^2-|h_1h_2|^2), \\
B_4 &= \frac{1}{2\Gamma\Gamma_2}\left[2\Im(g_1h_1^*g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)-(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)\right] \\
&- \frac{1}{2\Gamma\Gamma_1}\left[2\Im(g_1h_1^*g_2h_2^*)+(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)-(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\right] \\
&- \frac{\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}\left[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
&+ \frac{2}{\Gamma^2+\Delta m^2}\Im(g_1h_1^*g_2h_2^*), \\
C_0 &= A_0, C_3 = A_3, C_4 = A_4, \\
C_1 &= \frac{1}{2\Gamma_1^2}\left[|h_1h_2|^2-|g_1g_2|^2-(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
&- \frac{1}{2\Gamma_2^2}\left[|h_1h_2|^2-|g_1g_2|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
&- \frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}\left[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\right] \\
&+ \frac{\Gamma_1+\Gamma}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2(|g_1|^2-|h_1|^2)+\frac{2\Delta m}{\Gamma_1[(\Gamma+\Gamma_1)^2+\Delta m^2]}|g_2+h_2|^2\Im(g_1h_1^*) \\
&- \frac{\Gamma_2+\Gamma}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2(|g_1|^2-|h_1|^2)-\frac{2\Delta m}{\Gamma_2[(\Gamma+\Gamma_2)^2+\Delta m^2]}|g_2-h_2|^2\Im(g_1h_1^*) \\
&+ \frac{\Gamma(\Gamma_1+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2(|g_2|^2-|h_2|^2) \\
&- \frac{\Gamma(\Gamma_2+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2(|g_2|^2-|h_2|^2) \\
&- \frac{2\Delta m(\Gamma_2+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2\Im(g_2h_2^*) \\
&+ \frac{2\Delta m(\Gamma_1+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2\Im(g_2h_2^*) \\
&- \frac{4\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}\Im(g_1h_1^*g_2h_2^*),
\end{aligned}
$$

$$
\begin{aligned}
C_2 = {} & \frac{1}{2\Gamma_1^2}\bigl[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)+2\Im(g_1h_1^*g_2h_2^*)\bigr] \\
& + \frac{1}{2\Gamma_2^2}\bigl[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)-2\Im(g_1h_1^*g_2h_2^*)\bigr] \\
& + \frac{\Gamma^2-\Delta m^2}{(\Gamma^2+\Delta m^2)^2}\bigl[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\bigr] \\
& - \frac{2(\Gamma+\Gamma_1)}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2\Im(g_1h_1^*) - \frac{2(\Gamma_2+\Gamma)}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2\Im(g_1h_1^*) \\
& + \frac{\Delta m(\Gamma_1+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2(|g_2|^2-|h_2|^2) - \frac{2\Gamma\Delta m}{(\Gamma^2+\Delta m^2)^2}(|g_1g_2|^2-|h_1h_2|^2) \\
& + \frac{\Delta m(\Gamma_2+2\Gamma)}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2(|g_2|^2-|h_2|^2) \\
& - 2\frac{\Gamma(\Gamma_1+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_1+h_1|^2\Im(g_2h_2^*) \\
& - 2\frac{\Gamma(\Gamma_2+\Gamma)-\Delta m^2}{(\Gamma^2+\Delta m^2)[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_1-h_1|^2\Im(g_2h_2^*) \\
& + \frac{\Delta m}{\Gamma_2[(\Gamma_2+\Gamma)^2+\Delta m^2]}|g_2-h_2|^2(|g_1|^2-|h_1|^2) \\
& + \frac{\Delta m}{\Gamma_1[(\Gamma_1+\Gamma)^2+\Delta m^2]}|g_2+h_2|^2(|g_1|^2-|h_1|^2), \\
D_0 = {} & \frac{1}{8\Gamma\Gamma_1}|g_1-h_1|^2|g_2+h_2|^2 + \frac{1}{8\Gamma\Gamma_2}|g_1+h_1|^2|g_2-h_2|^2 \\
& - \frac{1}{4(\Gamma^2+\Delta m^2)}\bigl[(|g_1|^2-|h_1|^2)(|g_2|^2-|h_2|^2)+4\Im(g_1h_1^*)\Im(g_2h_2^*)\bigr] \\
& - \frac{\Delta m}{2\Gamma(\Gamma^2+\Delta m^2)}\bigl[(|g_1|^2-|h_1|^2)\Im(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Im(g_1h_1^*)\bigr], \\
D_1 = {} & \frac{1}{2\Gamma\Gamma_1}\bigl[|g_1h_2|^2-|g_2h_1|^2+(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\bigr] \\
& - \frac{1}{2\Gamma\Gamma_2}\bigl[|g_1h_2|^2-|g_2h_1|^2-(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)-(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\bigr] \\
& - \frac{1}{\Gamma^2+\Delta m^2}\bigl[(|g_1|^2-|h_1|^2)\Re(g_2h_2^*)+(|g_2|^2-|h_2|^2)\Re(g_1h_1^*)\bigr] \\
& - \frac{2\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}\Im(g_1h_1^*g_2h_2^*), \\
D_2 = {} & \frac{1}{2\Gamma\Gamma_1}\bigl[2\Im(g_1h_1^*g_2^*h_2)+(|h_1|^2+|g_1|^2)\Im(g_2h_2^*)+(|h_2|^2+|g_2|^2)\Im(g_1h_1^*)\bigr] \\
& - \frac{1}{2\Gamma\Gamma_2}\bigl[2\Im(g_1h_1^*g_2^*h_2)-(|h_1|^2+|g_1|^2)\Im(g_2h_2^*)-(|h_2|^2+|g_2|^2)\Im(g_1h_1^*)\bigr] \\
& - \frac{1}{\Gamma^2+\Delta m^2}\bigl[(|g_1|^2+|h_1|^2)\Im(g_2h_2^*)+(|g_2|^2+|h_2|^2)\Im(g_1h_1^*)\bigr] \\
& - \frac{\Delta m}{\Gamma(\Gamma^2+\Delta m^2)}(|g_1h_2|^2-|g_2h_1|^2),
\end{aligned}
$$

$$
\begin{aligned}
D_3 = {} & \frac{1}{2\Gamma\Gamma_1}\left[|g_1g_2|^2 - |h_1h_2|^2 + (|g_1|^2 - |h_1|^2)\Re(g_2h_2^*) - (|g_2|^2 - |h_2|^2)\Re(g_1h_1^*)\right] \\
& - \frac{1}{2\Gamma\Gamma_2}\left[|g_1g_2|^2 - |h_1h_2|^2 - (|g_1|^2 - |h_1|^2)\Re(g_2h_2^*) + (|g_2|^2 - |h_2|^2)\Re(g_1h_1^*)\right] \\
& + \frac{\Delta m}{\Gamma(\Gamma^2 + \Delta m^2)}\left[(|g_1|^2 + |h_1|^2)\Im(g_2h_2^*) - (|g_2|^2 + |h_2|^2)\Im(g_1h_1^*)\right] \\
& + \frac{1}{\Gamma^2 + \Delta m^2}(|g_1g_2|^2 - |h_1h_2|^2), \\
D_4 = {} & \frac{1}{2\Gamma\Gamma_1}\left[2\Im(g_1h_1^*g_2^*h_2) - (|h_1|^2 + |g_1|^2)\Im(g_2h_2^*) + (|h_2|^2 + |g_2|^2)\Im(g_1h_1^*)\right] \\
& - \frac{1}{2\Gamma\Gamma_2}\left[2\Im(g_1h_1^*g_2^*h_2) + (|h_1|^2 + |g_1|^2)\Im(g_2h_2^*) - (|h_2|^2 + |g_2|^2)\Im(g_1h_1^*)\right] \\
& + \frac{\Delta m}{\Gamma(\Gamma^2 + \Delta m^2)}\left[(|g_1|^2 - |h_1|^2)\Re(g_2h_2^*) - (|g_2|^2 - |h_2|^2)\Re(g_1h_1^*)\right] \\
& + \frac{2}{\Gamma^2 + \Delta m^2}\Im(g_1h_1^*g_2h_2^*).
\end{aligned}
$$

---

[1] A. Di Domenico (KLOE Collaboration), J. Phys. Conf. Ser. **171**, 012008 (2009).

[2] A. Di Domenico (KLOE Collaboration), Found. Phys. **40**, 852 (2010).

[3] G. Amelino-Camelia *et al.*, Europhys. J. C **68**, 619 (2010).

[4] For early consideration of EPR-correlated kaons, see M. Goldhaber, T. D. Lee and C. N. Yang, Phys. Rev. **112**, 1796 (1958); D. R. Inglis, Rev. Mod. Phys. **33**, 1 (1961); T. B. Day, Phys. Rev. **121**, 1204 (1961); H. J. Lipkin, *ibid* **176**, 1715 (1968).

[5] J. Bernabéu, F. J. Botella and J. Roldán, Phys. Lett. B **211**, 226 (1980); I. Dunietz, J. Hauser and J. L. Rosner, Phys. Rev. D **35**, 2166 (1987).

[6] C. Buchanan, R. Cousins, C. Dib, R.D. Peccei, J. Quackenbush, Phys.Rev. D **45**, 4088 (1992).

[7] G. D'Ambrosio, G. Isidori and A. Pugliese, hep-ph/9411389.

[8] J. Bernabéu, N. E. Mavromatos, and J. Papavassiliou, Phys. Rev. Lett. **92**, 131601 (2004); E. Álvarez and J. Bernabéu, JHEP **11**, 087 (2006); N. E. Mavromatos, Found. Phys. **40**, 917 (2010).

[9] K. R. S. Balaji, W. Horn and E. A. Paschos, Phys. Rev. D **68**, 076004 (2003).

[10] Y. Shi, Euro. Phys. J. C **72**, 1907 (2012).

[11] A. F. Falk and A. A. Petrov, Phys. Rev. Lett. **85**, 252 (2000); D. Atwood and A. A. Petrov, Phys. Rev. Lett. **71**, 054032 (2005); D. Atwood and A. Soni, Phys. Rev. D **82**, 036003 (2010).

[12] O. Kittle and A. Pilaftsis, arXiv:1108.3314.

[13] K. C. Chou, W. F. Palmer, E. A. Paschos, Y. L. Wu, Eur. Phys. J. C. **16**, 279-287 (2000).

[14] R. Louvot *et al.* (CLEO Collaboration), Phys. Rev. Lett. 102, 021801 (2009).

[15] A. Apostolakis (CPLEAR Collaboration), Phys. Lett B **422**, 339 (1998).